



ABOUT LUNG CANCER

Lung cancer occurs when cells in the lungs undergo abnormal changes in their DNA, leading to uncontrolled growth and the formation of a tumor. Over time, these cancerous cells can invade nearby tissues and potentially spread to other parts of the body through the bloodstream or lymphatic system.

Prevalence of Lung Cancer:

- Lung cancer is considered the primary cause of cancer-related deaths in both men and women in the United States.
- In 2018, 234,030 new instances of lung cancer were identified, constituting 13% of all cancer diagnoses.
- Lung cancer primarily affects older individuals. In 2015, 86% of individuals living with lung cancer were aged 60 or older.
- More men are diagnosed with lung cancer annually, while a greater number of women are living with the condition.
- Over 50% of individuals diagnosed with lung cancer do not survive beyond one year after receiving their diagnosis.





TYPES OF LUNG CANCER

Lung cancer comprises two primary categories: Non-small cell lung cancer and small cell lung cancer.

1. **Non-small-cell lung cancer (NSCLC):** This is the most common of the two types, accounting for about 85% of all lung cancer cases. NSCLC includes several subtypes, such as adenocarcinoma, squamous cell carcinoma, and large-cell carcinoma.
2. **Small-cell lung cancer (SCLC):** This type is less common, but tends to grow and spread more rapidly than NSCLC. It is often associated with a history of smoking.

Diagnosis, Treatment, and Prevention

Lung cancer is often diagnosed at an advanced stage because symptoms may not appear until the disease has progressed. Common symptoms can include a persistent cough, chest pain, coughing up blood, shortness of breath, unexplained weight loss, and fatigue.

The most significant risk factor for lung cancer is tobacco smoking, including cigarettes, cigars, and pipes. Environmental exposure to things like air pollution, secondhand smoke, and radon, as well as occupational exposure to carcinogens like asbestos, can also increase the risk. Additionally, genetic factors and family history may play a role in some cases.

Treatment for lung cancer depends on the type, stage, and overall health of the patient. It may include surgery, radiation therapy, chemotherapy, targeted therapy, immunotherapy, or a combination of these approaches. Early detection through screening, such as low-dose CT scans for high-risk individuals, can improve the chances of successful treatment.

Prevention of lung cancer primarily involves avoiding tobacco use and exposure to secondhand smoke, as well as minimizing exposure to environmental carcinogens. At-risk individuals should discuss their situation with healthcare professionals and undergo appropriate screening and risk-reduction measures.

MODIFIABLE RISK FACTORS

Modifiable risk factors for lung cancer are those factors that individuals have some control over and can potentially change or reduce their risk of developing the disease. The most significant modifiable risk factor for lung cancer is cigarette smoking, which accounts for the majority of lung cancer cases. **Here are the primary modifiable risk factors for lung cancer:**




- **Tobacco Smoking:** Smoking tobacco, whether in the form of cigarettes, cigars, or pipes, is the leading cause of lung cancer. It is responsible for a vast majority of lung cancer cases worldwide. Quitting smoking, or never starting in the first place, is the most effective way to reduce the risk.
- **Secondhand Smoke:** Exposure to secondhand smoke, also known as passive smoking, can increase the risk of lung cancer, especially in nonsmokers. Avoiding environments where smoking is prevalent and ensuring a smoke-free home and workplace can reduce this risk.
- **Vaping and E-Cigarettes:** While e-cigarettes or vaping products may be perceived as less harmful than traditional cigarettes, they are not without risk. Vaping can expose individuals to harmful chemicals and carcinogens, potentially increasing the risk of lung cancer. It's essential to be cautious and informed about the risks associated with these products.
- **Radon Exposure:** Radon is a naturally occurring radioactive gas that can seep into homes and buildings from the ground. Prolonged exposure to high radon levels can increase the risk of lung cancer. Testing homes for radon and taking measures to mitigate radon levels, if necessary, can reduce this risk.
- **Occupational Exposures:** Some workplaces, such as construction sites and industrial settings, may expose workers to carcinogens like asbestos, arsenic, and diesel exhaust. Proper safety measures, protective equipment, and adherence to workplace regulations can help reduce these exposures.
- **Air Pollution:** Long-term exposure to outdoor air pollution, particularly in areas with high fine particulate matter (PM2.5) and other pollutants, can contribute to lung cancer risk. Reducing exposure by staying indoors during poor air quality days and supporting clean air initiatives can help mitigate this risk.
- **Diet and Nutrition:** While the direct link between diet and lung cancer is complex, a healthy diet rich in fruits and vegetables may have a protective effect. Research indicates that an increased intake of fruits and vegetables abundant in antioxidants is associated with a decreased risk of certain cancers, including lung cancer.
- **Physical Activity:** There is some indication that engaging in regular physical activity may reduce the likelihood of developing lung cancer.

NON-MODIFIABLE RISK FACTORS

Non-modifiable risk factors for lung cancer are factors that individuals cannot change or control. These risk factors are typically related to a person's genetics, age, and certain environmental exposures that may have occurred in the past. While you cannot modify these risk factors, understanding them can still be important for assessing your overall risk of developing lung cancer. Here are some non-modifiable risk factors for lung cancer:

- **Age:** Lung cancer risk increases with age, and it is most commonly diagnosed in individuals over the age of 65. The risk tends to rise significantly with advancing age.
- **Family History:** Having a family history of lung cancer may increase your risk, as it may suggest a genetic predisposition or shared environmental exposures within the family. However, most cases of lung cancer occur in individuals with no family history of the disease.
- **Genetics:** Certain genetic mutations or variations can increase the susceptibility to lung cancer. For example, mutations in genes like EGFR (epidermal growth factor receptor) and ALK (anaplastic lymphoma kinase) are associated with a higher risk of specific types of non-small-cell lung cancer (NSCLC).
- **Certain Lung Diseases:** Some pre-existing lung conditions, such as chronic obstructive pulmonary disease (COPD), pulmonary fibrosis, and tuberculosis, may raise the risk of developing lung cancer.
- **Race and Ethnicity:** Lung cancer rates can vary among different racial and ethnic groups. For example, in the United States, African American men have a higher lung cancer incidence rate compared to white men, although the reasons for these disparities are multifactorial and not fully understood.

If you have concerns about your risk of lung cancer, especially if you have non-modifiable risk factors, discuss them with a healthcare professional. They can guide appropriate screening and risk reduction strategies based on your circumstances.



LUNG CANCER SCREENING TESTS



Lung cancer screening tests are medical examinations and procedures designed to detect lung cancer at an early stage, typically before symptoms become apparent. Early detection can improve the chances of successful treatment and increase survival rates.

Lung cancer screening is recommended primarily for individuals who are at a higher risk of developing the disease, particularly those with a history of smoking.

The most common lung cancer screening test is low-dose computed tomography (LDCT). Here's an overview of lung cancer screening tests:

Low-Dose Computed Tomography (LDCT)

- LDCT is the most widely-used and recommended screening test for lung cancer.
- It involves a series of X-ray images taken from various angles to create detailed cross-sectional images of the lungs.
- The "low-dose" aspect refers to the use of a lower radiation dose compared to standard CT scans, reducing the risk of radiation exposure.
- LDCT can detect small lung nodules, including those that may be early-stage lung cancers.
- The National Lung Screening Trial (NLST) in the United States demonstrated that LDCT screening reduced lung cancer mortality in high-risk individuals compared to chest X-rays.

Chest X-rays:

- Chest X-rays were historically used for lung cancer screening, but they are less effective than LDCT at detecting early-stage lung cancer.

- They are not recommended as a primary screening tool but may be used in some cases or in combination with other tests.

Sputum Cytology:

Sputum cytology involves analyzing a sample of mucus coughed up from the lungs to look for abnormal cells.

It is less sensitive and specific than LDCT and is not commonly used for routine lung cancer screening.

Screening Criteria:

It's important to note that lung cancer screening is typically recommended for individuals who meet specific criteria and are at higher risk. Common eligibility criteria for lung cancer screening in many guidelines include:

- **Age:** Usually between 55 and 80 years old.
- **Smoking History:** A significant history of smoking, often defined as a certain number of pack-years (e.g., 30 pack-years or more).
- **Current or Former Smokers:** Current smokers or individuals who have quit smoking within the past 15 years.
- **Overall Health:** Generally good overall health to tolerate potential follow-up procedures or treatments if abnormalities are found.

The decision to undergo lung cancer screening should be made in consultation with a healthcare provider, who can assess an individual's risk factors and determine if screening is appropriate.

Additionally, healthcare providers can discuss the benefits and potential risks of screening, such as false-positive results and the need for further testing or procedures.

Lung cancer screening programs are most effective when combined with smoking cessation efforts and discussions about risk reduction. Quitting smoking is the most significant step an individual can take to reduce their risk of lung cancer, regardless of their eligibility for screening.

REFERENCES

1. American Lung Association. (n.d.). Lung.org. <https://www.lung.org>
2. Jonas, D. E., Reuland, D. S., Reddy, S. M., Nagle, M., Clark, S. D., Weber, R. P., ... & Harris, R. P. (2021). Screening for lung cancer with low-dose computed tomography: Updated evidence report and systematic review for the US Preventive Services Task Force. **Jama**, 325(10), 971-987.
3. National Cancer Institute. (n.d.). Lung Cancer. National Cancer Institute. <https://www.cancer.gov/types/lung>
4. Siegel, R. L., Miller, K. D., & Jemal, A. (2018). Cancer Statistics, 2018. **CA: A Cancer Journal for Clinicians**, 68(1), 7-30. <https://doi.org/10.3322/caac.21442>
5. U.S. National Institute of Health, National Cancer Institute. (2018). SEER Cancer Statistics Review, 1975-2015.
6. Yang, J., Qian, S., Na, X., & Zhao, A. (2023). Association between Dietary and Supplemental Antioxidants Intake and Lung Cancer Risk: Evidence from a Cancer Screening Trial. **Antioxidants (Basel, Switzerland)**, 12(2), 338. <https://doi.org/10.3390/antiox12020338>

